

We claim:

1. A hydrophilic matrix formulation suitable for once-a-day administration comprising:
  - a. divalproex sodium, and;
  - 5 b. said divalproex sodium is in admixture with a sufficient quantity of a pharmaceutically acceptable polymer, so that said formulation exhibits the following in-vitro dissolution profile, when measured in a type 2 dissolution apparatus (paddle) at 100 rpm, at a temperature of  $37 \pm 0.5$  C, in 500ml of 0.1N HCl for 45 minutes, followed by 900ml of 0.05M phosphate buffer containing 75.  
10 mM sodium laurel sulfate ( pH5.5) for the remainder of the testing period:
    - i. no more than about 30 % of total valproate is released after 3 hours of measurement in said apparatus;
    - ii. from about 40 to about 70% of total valproate is released after 9 hours of measurement in said apparatus;
    - 15 iii. from about 55 to about 95% of total valproate is released after 12 hour of measurement in said apparatus, and;
    - iv. not less than 85% of total valproate is released after 18 hours of measurement in said apparatus.
2. The formulation according to claim 1 in which said formulation exhibits the  
20 following in-vitro dissolution profile:
  - i. from about 15% to about 30% of total valproate is released after 3 hours of measurement in said apparatus;
  - ii. from about 40% to about 70% of total valproate is released after 9 hours of measurement in said apparatus;
  - 25 iii. from about 55% to about 90% of total valproate is released after 12 hours of measurement in said apparatus, and;
  - iv. not less than 88% of total valproate is released after 18 hours of measurement in said apparatus.

3. The formulation according to claim 1 in which said formulation exhibits the following in-vitro dissolution profile:
  - i. from about 15% to about 27% of total valproate is released after 3 hours of measurement in said apparatus;
  - 5 ii. from about 44% to about 69% of total valproate is released after 9 hours of measurement in said apparatus;
  - iii. from about 59% to about 90% of total valproate is released after 12 hours of measurement in said apparatus, and ;
  - 10 iv. not less than 88% of total valproate is released after 18 hours of measurement in said apparatus.
4. The formulation according to claim 1 in which said divalproex sodium is present in the amount of from about 40 to about 80w/w% based upon the total weight of the formulation.
5. The formulation according to claim 3 in which said polymer is a water soluble  
15 hydrophilic polymer is selected from the group consisting of polyvinylpyrrolidine, hydroxypropyl cellulose, hydroxypropylmethyl cellulose, methyl cellulose, vinyl acid copolymers, methacrylic acid copolymers, maleic anhydride/methyl vinyl ether copolymers and mixtures thereof.
6. The formulation according to claim 5 in which said divalproex sodium is present in the  
20 amount of from about 45 to about 65 w/w%, based upon the total weight of the formulation.
7. The formulation according to claim 6 in which said polymer is present in the amount of from about 20 to about 50 w/w%, based upon the total weight of the formulation.
8. The formulation according to claim 7 which further comprises one or more  
25 pharmaceutically acceptable excipients.
9. A method for treating migraine comprising administering a formulation according to claim 1 to a patient in need thereof.
10. A method for treating epilepsy comprising administering a formulation according to claim 1 to a patient in need thereof.

11. A method for treating bipolar disorders comprising administering a formulation according to claim 1 to a patient in need thereof.
12. The formulation according to claim 1, which when ingested orally produces a  $C_{\max}$  that is statistically significantly lower than the  $C_{\max}$  produced by a delayed release divalproex sodium tablet, when each is determined at steady state in a fasting population.
13. The formulation according to claim 12 which:
- a) produces a  $C_{\min}$  that is not statistically significantly different from the  $C_{\min}$  produced by said delayed release divalproex sodium tablet, when each is determined at steady state in a fasting population, and;
  - b) said formulation produces an AUC value that is equivalent to the AUC value generated by said divalproex sodium delayed release tablet, when each is determined at steady state in a fasting population.
14. A hydrophilic matrix formulation suitable for once-a-day administration comprising:
- a. divalproex sodium, and;
  - b. said divalproex sodium is in admixture with a sufficient quantity of a pharmaceutically acceptable polymer, so that said formulation exhibits the following in-vitro dissolution profile, when measured in a type 2 dissolution apparatus (paddle) at 100 rpm, at a temperature of  $37 \pm 0.5$  C, in 500ml of 0.1N HCl for 45 minutes; followed by 900ml of 0.05M phosphate buffer containing 75 mM sodium laurel sulfate (pH5.5) for the remainder of the testing period:
    - i. from about 15% to about 27% of total valproate is released after 3 hours of measurement in said apparatus;
    - ii. from about 44% to about 69% of total valproate is released after 9 hours of measurement in said apparatus;
    - iii. from about 59% to about 90% of total valproate is released after 12 hours of measurement in said apparatus, and;
    - iv. not less than 88% of total valproate is released after 18 hours of measurement in said apparatus.

15. The formulation according to claim 14, which when ingested orally produces a  $C_{\max}$  that is statistically significantly lower than the  $C_{\max}$  produced by a delayed release divalproex sodium tablet, when each is determined at steady state in a fasting population.

5 16. A hydrophilic matrix formulation suitable for once-a-day administration comprising:

a) a valproate compound, and;

b) said valproate compound is in admixture with a sufficient quantity of a pharmaceutically acceptable polymer, so that said formulation exhibits the following in-vitro dissolution profile, when measured in a type 2 dissolution apparatus (paddle) at 100 rpm, at a temperature of  $37 \pm 0.5^{\circ}\text{C}$ , in 500ml of 0.1N HCl for 45 minutes, followed by 900ml of 0.05M phosphate buffer containing 75 mM sodium laurel sulfate, pH5.5, for the remainder of the testing period:

- 10
- 15
- 20
- i. no more than about 30 % of total valproate is released after 3 hours of measurement in said apparatus;
  - ii. from about 40 to about 70% of total valproate is released after 9 hours of measurement in said apparatus;
  - iii. from about 55 to about 95% of total valproate is released after 12 hour of measurement in said apparatus, and;
  - iv. not less than 85% of total valproate is released after 18 hours of measurement in said apparatus.

17. The formulation according to claim 1 in which said formulation exhibits the following in-vitro dissolution pattern:

- 25
- 30
- i. from about 15% to about 30% of total valproate is released after 3 hours of measurement in said apparatus;
  - ii. from about 40% to about 70% of total valproate is released after 9 hours of measurement in said apparatus
  - iii. from about 55% to about 90% of total valproate is released after 12 hours of measurement in said apparatus

- iv. not less than 88% of total valproate is released after 18 hours of measurement in said apparatus.

18. The formulation according to claim 1 in which said formulation exhibits the following in-vitro dissolution pattern:

- 5           i. from about 15% to about 27% of total valproate is released after 3 hours of measurement in said apparatus;
- ii. from about 44% to about 69% of total valproate is released after 9 hours of measurement in said apparatus
- 10          iii. from about 59% to about 90% of total valproate is released after 12 hours of measurement in said apparatus
- iv. not less than 88% of total valproate is released after 18 hours of measurement in said apparatus.